

NEUROPATHIC PAIN

CJ Cullinan RN BSN CRRN MHSA

Craig Rehabilitation Hospital
Englewood CO

cjcullinan@msn.com

Conditions with NP

- Diabetic peripheral neuropathy
- Post herpetic neuralgia
- HIV neuropathy
- Spinal cord injury
- Neuropathic cancer pain
- Phantom limb pain
- post-mastectomy pain
- Guillian-Barre ´ syndrome
- Chemotherapy-induced neuropathy
- Complex regional pain syndrome, types I & II
- Vitamin B12 Deficiency
- Spinal stenosis
- Lupus
- Multiple sclerosis
- Fibromyalgia
- Lead ingestion
- Traumatic injury: motor vehicle accident or gunshot
- Failed back surgery syndrome

Etcetera, etcetera, etcetera.....

2

Numbers/Demographics

Estimated US Population 304,500,000
Medscape estimates 3% of our population have NP!

- ◆ Those with diabetes, 25% develop NP
- ◆ Of the 1 million annual cases of Shingles, 20% develop post-herpetic neuralgia
- ◆ Most common reason patients seek medical care is for pain
 - 20% for persistent chronic pain
- ◆ **Specialists theorize that most chronic pain is NP**

(Finnerup, 2007; CDC, 2008)

Neuropathic Pain (NP) Definition

“Initiated or caused by
a primary lesion or dysfunction
in the nervous system”

- The International Association for the Study of Pain (IASP)

Further clarifications:

- ◆ Make a distinction between nociceptive and NP
- ◆ Differentiate from peripherally-generated and centrally-generated NP
- ◆ Note structure and process differences - neuroplasticity

4

Symptoms

- ◆ **Allodynia** – pain caused by something that is generally non-painful (light touch)
- ◆ **Hyperalgesia** - heightened response to normally painful stimulus
- ◆ **Hyperesthesias** – an exaggerated response to touch, such as to bed sheets
- ◆ **Hyperpathy** – persistent pain even after the cause of pain has been removed
- ◆ **Paresthesias and dyesthesias** – abnormal and unpleasant sensations that are described as tingling, pins and needles
- ◆ **Abherent algia** the pt feels requires continual movement, touch
- ◆ **Worse at night**
- ◆ **Worse after weight bearing**

(Svendsen, 2006; Parry/MacCarberg, 2007)


5

Pathophysiology

Basic nociceptive pain includes:

- ◆ **Transduction at the nociceptors**
 - Thermal, Chemical, or Mechanical stimuli
- ◆ **Transmission via C and A fibers**
- ◆ **Modulation**
 - Dorsal horn
 - Chemical receptors on the axon membrane
 - ◆ Opiate, GABA, Bradykinin, Histamine, Serotonin, Capsaicin
- ◆ **Perception**


(Abrams, 2007)



Abherent Neuroprocessing

- Sodium channels of unusual density produce persistent spontaneous discharge
- Calcium channels increase Substance P, glutamate, and norepinephrine
- Aberrant processing continues in PNS/CNS
- Receptor field size increases

(Tremont-Lucats, 2006, Hornick, 2007)



AND THAT MEANS???

The nerves are excited!!!!!!!!!!!!

◆ GOALS FOR NP TREATMENT

- Restore inhibitory neurotransmission – Sodium channels
- Modulate Calcium channels
- Block peripheral sensitization
- Block central sensitization
- "Retrain the nervous system"

(Hornick, 2007)

10

Assessment Tools

- ◆ Pain Quality Assessment Scale (PQAS) (2008)
- ◆ Self-Administered Leeds Assessment of Neuropathic Symptoms and Signs (S-LANSS) (2005)
- ◆ Douleur Neuropathique 4 questions questionnaire (DN4) (2004)
- ◆ 12 item Neuropathic Pain Questionnaire (NPQ) (2003)
- ◆ The Neuropathic Pain Scale (1996)
- ◆ Quantitative sensory testing (QST) (1994)

11

Physical Assessment

- ◆ Pain location
- ◆ Distribution
- ◆ Intensity
- ◆ Quality
- ◆ Time course
- ◆ Underlying disease
- ◆ Area of possible nervous system lesion

(Jensen, 2003)

12

Lab Tests

- ◆ CBC with Diff
- ◆ Sed rate
- ◆ Chem profile
- ◆ Thyroid
- ◆ B12
- ◆ Folate
- ◆ Fasting blood sugar
- ◆ Gyclo Hgb
- ◆ Protein electrophoresis
- ◆ Lyme titers
- ◆ Hep B/C
- ◆ HIV
- ◆ Anti-nuclear
- ◆ Sjogren's titers
- ◆ Anitneutrophil cytoplasmic antibody
- ◆ Rheumatoid factor
- ◆ Cryoglobulins
- ◆ Anti-sulfide antibody titers
- ◆ Anti-HU titers
- ◆ Heavy metal – serum/urine screens
- ◆ CSF for demylinating diseases & meningeal carcinomatosis
- ◆ Biopsies Dx for vasculitis, amyloidosis, sarcoidosis, etc.

(Beth Israel Medical Center, 2008)

13

McHUMOR.com by T. McCracken



"I'd say the nerve pain you're having in your jaw is due to an arrow through it, but perhaps you'd like to get a 2nd opinion from a dentist."


14

CONSIDERATIONS FOR TX

- ◆ Co-morbidities and complexities
- ◆ Potential adverse outcomes
- ◆ Potential drug interactions
- ◆ Costs associated with therapy
- ◆ Potential risks of medication abuse
- ◆ Individual variation
- ◆ Multimodalities
 - Polypharmacy
 - Can include pharmacological-nonpharmacological combinations

(Dworkin, 2007)


15



TREATMENT NOTE:

Success is a pain reduction of 40-60%.

(Parry, 2006)



FIRST LINE MEDICATIONS

- Calcium Channel Blockers/Modulators

- ◆ **Gabapentin (Neurontin®)**
 - + worsens cognitive or gait issues for elderly
- ◆ **Pregabalin (Lyrica®)**
 - + concern with renal impairment
 - + weight gain

(Dworkin, 2007)
19

FIRST LINE Continued

- Tricyclic Antidepressant (TCA)

- ◆ **Nortriptyline/Pamelor®**
- ◆ **Desipramine/Norpramin®**

- Monitor EKG
- Strong drug interactions with SSRIs
- Not recommended:
amitriptyline (Elavil®) & imipramine (Tofranil®)

(Dworkin, 2007)
20

FIRST LINE Continued

- SSNRI - Selective serotonin & norepinephrine reuptake inhibitor

- ◆ **Duloxetine (Cymbalta®)** antidepressant
 - Contraindicated: uncontrolled glaucoma, end-stage renal
 - Monitor BP, suicide ideations
- ◆ **Venlafaxine (Effexor®)** antidepressant
 - Monitor EKG

(Finnerup, 2007)
21

SECOND AND THIRD LINE

◆ Opioid analgesics

- Proven less sensitive, but assists with intense pain

◆ Tramadol (Ultram®)

- Don't use with SSRI, possible serotonin syndrome

◆ IV morphine

◆ IV lidocaine

- dissociative anesthetic

(Dworkin, 2007)
22

Other Medications Used

◆ Methadone

- Looooooooooooooooong half life
- Contraindicated with liver dysfunction
- When changing from another opioid to methadone, reduce dose 75-95% of expected equianalgesic

◆ Buprenorphine

(Buprenex®, Subutex®)

- systemic potency 20-40 times > morphine
- long acting, low CNS effects, less constipation
- Used transdermally for up to 96hr
- Suboxone® is a combo of buprenorphine and naloxone

(Likar, 2006; Kress, 2008)
23

Other Medications Used Continued

- Calcium Channel Blockers/Modulators

◆ Topiramate (Topamax®)

- Studies show noneffective, however...

◆ Lamotrigine (Lamictal®)

- Black box warning re: Stevens-Johnson

◆ Benzodiazepines

- Clonazepam (Klonopin®)
- Valium®

24

Other Medications Used Continued

◆ **NMDA** (*N*-methyl-D-aspartic acid) antagonists

- Dextromethorphan
- Ketamine
- Nemantine (Namenda®)
- Riluzole (Rilutek®)
- Amantadine

◆ **Alpha-2-adrenergic agonists**

- Clonidine
 - ◆ PO, topical, intrathecal

25

Other Medications Used Continued

◆ **Mexiletine po** (Mexitil®)

- Oral lidocaine
- Limited use r/t cardiac effects, nausea

◆ **Cholecystokinin-B antagonist** (Proglumide)

◆ **Calcitonin** Osteoporosis

◆ **Flecainide** (Tambocor ®)

◆ **Baclofen** Antispasmodic


◆ **Vitamin D supplementation**

◆ **Levetiracetam** (Keppra®) Anticonvulsant

26



What NOT to use for Neuropathic Pain



(Parry, 2007- Copyright © 2006 Dannemiller Memorial Educational Foundation - via Medscape)

27

TOPICALS AND PATCHES

◆ Patches


- Lidocaine – Lidoderm®
 - ◆ Patch and Gel
- Clonidine
- Diclofenac – Flector®, Voltaren®
- Buprenorphine - BuTrans patch®
- Sufentanil – Transdur®
- Bupivacaine - Eladur®

(Lyda, 2008)

◆ Alternatives

- Camphor (*Cinnamomum camphora*)
- Capsicum (*Capsicum* species)
- ◆ Counter irritants, likely safe

(Natural Medicine, 2008)
28





Interventional


◆ Spinal Cord Stimulators – Implanted

- Consider when polypharmacy failed
- The pathway is intact
- Successful external stimulator trial over 5-7 days

(Varrassi, 2005)
30







◆ **Tunneled Epidural Catheter**

◆ **Motor Cortex Stimulator** –
neurosurgical craniotomy

◆ **Peripheral stimulation Unit** –
implanted

◆ **Phenol sympathectomy**
– No positive effects

(Varrassi, 2005)

34

◆ **Implanted Intrathecal Catheter**

When to use is similar to
stimulators:

- Polypharmacy failed
- Side Effects too intense

- **Clonidine**
• usually in combination with
other anesthetics or opioids

- **Ziconotide (Prialt®)**
calcium channel blocker



Madaris, 2008

35

Complementary & Alternative

◆ **Acupuncture**
– Various types of acupuncture

◆ **Aromatherapy**
Massage, Baths, Inhalations

– **Essential Oils**
Juniper, V-6, peppermint, NeuroGen®,
chamomile, sandalwood

– **Patchouli incense**

◆ **Acetyl-L-carnitine**

36

CAM Continued

- ◆ **Alpha-lipoic acid**

- ◆ **St. John's Wort** (*Hypericum perforatum*)
 - Check for drug-herb interactions

- ◆ **Cannabinoids**

- Dronabinol (Marinol®) provides significant pain relief (Finnerup, 2007)

- ◆ **CONSIDERED UNSAFE**

- L-tryptophan
- 5-HTP (Natural Medicine, 2008)

37

TREATMENT SUPPORT

- ◆ Stress reduction
- ◆ Good sleep hygiene
- ◆ Diet with protein/amino acids
- ◆ Physical activity
 - Walking, swimming
 - Physical Therapy
 - ◆ Assess for balance
 - ◆ Gentle Desensitization
 - ◆ Progress to flexibility and isometric strengthening exercises (AMA, 2007)

38

Is Neuropathic Pain Preventable?

- ◆ **Peri-operative with gabapentin or pregabalin**
 - ◆ IASP states 1200mg night before plus 5 days post-op
- ◆ **Tight Diabetic Glycemic Control**

(Campbell, 2006; AMA, 2007)

39

Continue the Prevention

◆ Treat Pain Consistently

- Repetitive noxious stimulation present
- "Wind up" - Progressive increase in # of action potentials
- Treat pain when it happens, don't let it become chronic and/or neuropathic

◆ Herpes Zoster Vaccine

- Of those receiving the vaccine yet still acquiring shingles, 67% decrease in incidence of PHN

(CDC, 2008; Perry, 2006)

EVALUATION

ANY TREATMENT WILL TAKE TIME

- Achieve a steady state
- Titrate to 'effect and side effect'
- Reevaluate to increase/decrease
- Keep documentation about the trials - successes and failures
 - ◆ "What worked" "For how long" "What didn't work"
- Encourage the patient to keep their own records
 - ◆ **THEIR involvement is a MUST!**

41



42

General Rule for Treatment

- ◆ If effective, continue.
- ◆ If partially effective, consider multimodalities.
- ◆ If not effective, change your intervention. Try something else.

(Dworkin, 2007)

43

Thank you!

CJ Cullinan RN BSN CRRN MHSA
cjcullinan@msn.com

44
